

N-ARY QUASIGROUPS OF ORDER 4

DENIS S. KROTOV* AND VLADIMIR N. POTAPOV†

Abstract. We characterize the set of all n -ary quasigroups of order 4: every n -ary quasigroup of order 4 is permutably reducible or semilinear. Permutable reducibility means that an n -ary quasigroup can be represented as a composition of k -ary and $(n - k + 1)$ -ary quasigroups for some k from 2 to $n - 1$, where the order of arguments in the representation can differ from the original order. The set of semilinear n -ary quasigroups has a characterization in terms of Boolean functions.

Key words. Latin hypercube, n -ary quasigroup, reducibility

AMS subject classifications. 05B15, 20N05, 20N15, 94B25

1. Introduction. An algebraic system consisting of a finite set Σ of cardinality $|\Sigma| = q$ and an n -ary operation $f : \Sigma^n \rightarrow \Sigma$ uniquely invertible in each place is called an n -ary quasigroup of order q . The function f can also be referred to as an n -ary quasigroup of order q or, for short, an n -quasigroup. The value array of an n -quasigroup of order q is known as a latin n -cube of order q (if $n = 2$, a latin square). Furthermore, there is a one-to-one correspondence between the n -quasigroups and the distance 2 MDS codes in Σ^{n+1} .

It is known that for every n there exist exactly two equivalent n -quasigroups of order 2 and $3 \cdot 2^n$ n -quasigroups of order 3, which constitute one isotopy class (see, e.g., [LM98]). So, 4 is the first order for which a rich class of n -quasigroups exists. On the other hand, this order is of special interest for different areas of mathematics close to information theory. For example,

- the class of 1-perfect codes in $\{0, 1\}^n$ of rank at most $n - \log_2(n + 1) + 2$ (the minimum rank is $n - \log_2(n + 1)$ for 1-perfect codes) is characterized in terms of n -quasigroups of order 4, see [AHS04] (so, our work completes this characterization);
- order 4 is the first order that is applicable for use in quasigroup stream ciphers;
- from n -quasigroups of order 4, n -quasigroups of other orders can be constructed, giving examples of n -quasigroups with nontrivial properties (see, e.g., [Kro08a]).

In this paper, we show that every n -quasigroup of order 4 is permutably reducible or semilinear. Permutable reducibility means that the n -quasigroup can be represented as a repetition-free composition of quasigroups of smaller arities where the ordering of the arguments in the representation can differ from the original (see Definition 2.4). Semilinearity (Definition 2.5) means that the n -quasigroup can be obtained as a direct product of two n -quasigroups of order 2 modified by a Boolean function $\{0, 1\}^n \rightarrow \{0, 1\}$ (sometimes this construction is referred to as the wreath product construction, but we should remember that this does not agree with the concept of wreath product of groups).

In §2 we introduce main concepts and notations. In §3 we formulate the result (Theorem 3.1) and divide the proof into four subcases, Lemmas 3.2-3.5. Lemmas 3.4 and 3.5 are proved in §5 and §4, while Lemmas 3.2 and 3.3 follow from previous

*Sobolev Institute of Mathematics, prosp. Akademika Koptiyuga, 4, Novosibirsk, 630090, Russia (krotov@math.nsc.ru).

†Sobolev Institute of Mathematics, prosp. Akademika Koptiyuga, 4, Novosibirsk, 630090, Russia (vpotapov@math.nsc.ru).

papers.

2. Main definitions.

DEFINITION 2.1. An n -ary operation $Q : \Sigma^n \rightarrow \Sigma$, where Σ is a nonempty set, is called an n -ary quasigroup or n -quasigroup (of order $|\Sigma|$) if in the equality $z_0 = Q(z_1, \dots, z_n)$ knowledge of any n elements of $z_0, z_1, \dots, z_n$ uniquely specifies the remaining one [Bel72].

The definition is symmetric with respect to the variables $z_0, z_1, \dots, z_n$, and sometimes it is convenient to use a symmetric form for the relation $z_0 = Q(z_1, \dots, z_n)$. For this reason, we will denote by $Q\langle z_0, z_1, \dots, z_n \rangle$ the corresponding predicate, i. e., the characteristic function of this relation. (In coding theory, the set corresponding to this predicate is known as a distance 2 MDS code.)

Given $\bar{y} = (y_1, \dots, y_n)$, we denote

$$\bar{y}^{[i]}[x] \triangleq (y_1, \dots, y_{i-1}, x, y_{i+1}, \dots, y_n);$$

similarly, we define $\bar{y}^{[i_1, i_2, \dots, i_k]}[x_{i_1}, x_{i_2}, \dots, x_{i_k}]$.

DEFINITION 2.2. If we assign some fixed values to $l \in \{1, \dots, n\}$ variables in the predicate $Q\langle z_0, \dots, z_n \rangle$ then the $(n - l + 1)$ -ary predicate obtained corresponds to an $(n - l)$ -quasigroup. Such a quasigroup is called a retract or $(n - l)$ -retract of Q . If z_0 is not fixed, the retract is principal.

DEFINITION 2.3. By an isotopy we shall mean a collection of $n + 1$ permutations $\tau_i : \Sigma \rightarrow \Sigma$, $i \in \{0, 1, \dots, n\}$. n -Quasigroups f and g are called isotopic, if for some isotopy $\bar{\tau} = (\tau_0, \tau_1, \dots, \tau_n)$ we have $f(x_1, \dots, x_n) \equiv \tau_0^{-1}g(\tau_1x_1, \dots, \tau_nx_n)$, i. e., $f\langle x_0, x_1, \dots, x_n \rangle \equiv g\langle \tau_0x_0, \tau_1x_1, \dots, \tau_nx_n \rangle$.

DEFINITION 2.4. An n -quasigroup f is termed permutably reducible (in [PK06], the term “decomposable” was used) if there exist $m \in \{2, \dots, n - 1\}$, an $(n - m + 1)$ -quasigroup h , an m -quasigroup g , and a permutation $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ such that

$$f(x_1, \dots, x_n) \equiv h(g(x_{\sigma(1)}, \dots, x_{\sigma(m)}), x_{\sigma(m+1)}, \dots, x_{\sigma(n)})$$

(i. e., f is a composition of h and g). For short, we will omit the word “permutably” (with the exception of the main statements). If an n -quasigroup is not reducible, then it is irreducible. (In particular, all 2-quasigroups are irreducible.)

DEFINITION 2.5. We say that an n -quasigroup $f : \{0, 1, 2, 3\}^n \rightarrow \{0, 1, 2, 3\}$ is standardly semilinear if

$$f\langle \bar{x} \rangle \leq L\langle \bar{x} \rangle$$

where

$$L\langle x_0, \dots, x_n \rangle \triangleq l(x_0) \oplus \dots \oplus l(x_n) \oplus 1, \quad l(0) = l(1) = 0, \quad l(2) = l(3) = 1$$

($\leq$ means “ $\leq$ everywhere”; $\oplus$ means modulo-2 addition), see, e. g., Fig. 2.1.

An n -quasigroup of order 4 is called semilinear if it is isotopic to some standardly semilinear n -quasigroup.

The set of standardly semilinear n -quasigroups has a simple characterization:

PROPOSITION 2.6. The following relation is a bijection between the standardly semilinear n -quasigroups f and the Boolean functions $\lambda : \{0, 1\}^n \rightarrow \{0, 1\}$:

$$(2.1) \quad f\langle x_0, x_1, \dots, x_n \rangle \equiv L\langle x_0, x_1, \dots, x_n \rangle \cdot (x_0 \oplus x_1 \oplus \dots \oplus x_1 \oplus \lambda(l(x_1), \dots, l(x_n))).$$




FIGURE 2.1. A standardly semilinear 2-quasigroup and the function $L(\cdot)$

Proof. Consider an standardly semilinear n -quasigroup. Consider a set H consisting of 2^{n+2} points of $\{0, 1, 2, 3\}^{n+1}$ with fixed values $l(x_1), \dots, l(x_n)$.

The number of 1s of $f(\cdot)$ in H is 2^n (indeed, by the definition of an n -quasigroup, every n -tuple $x_1, \dots, x_n$ corresponds to exactly one 1). Moreover, since f is standardly semilinear, all these 1s belong to $H_1 \triangleq (\bar{x} \in H \mid L(\bar{x}) = 1)$. Since there are no two 1s that differ in only one coordinate, all these 1s simultaneously have either even or odd coordinate sum. In the even case define $\lambda(l(x_1), \dots, l(x_n)) = 1$; in the odd case, $= 0$. Then (2.1) is automatically true. $\square$

So, the number of the standardly semilinear n -quasigroups is 2^{2^n} . Multiplying by the number $3^{n+1}2$ of different functions isotopic to L , we obtain an approximate number of the semilinear n -quasigroups. The exact number is $3^{n+1}2^{2^n+1} - 8 \cdot 6^n$ [PK06, Theorem 1], where $-8 \cdot 6^n$ is explained by the fact that affine Boolean functions (and only affine, i.e., of type $\lambda(z_1, \dots, z_n) = b_0 \oplus b_1 z_1 \oplus \dots \oplus b_n z_n$, $b_i \in \{0, 1\}$) corresponds to n -quasigroups majorized by more than one isotope of L .

In the rest of the paper, unless otherwise stated, we consider only order-4 n -quasigroups over $\Sigma = \{0, 1, 2, 3\}$.

3. Main result. The main result is the following theorem.

THEOREM 3.1. *Every n -quasigroup of order 4 is permutably reducible or semilinear.*

The basic characteristic of an n -quasigroup f , which divides our proof into four subcases, is the maximum arity of its irreducible retract. Denote this value by $\kappa(f)$; then, $2 \leq \kappa(f) < n$. The line of reasoning in the proof of Theorem 3.1 is inductive, so we can assume that the irreducible retracts are semilinear.

LEMMA 3.2 (Case $\kappa = n - 1$, [PK06, Lemma 4]). *If an n -quasigroup f of order 4 has a semilinear $(n - 1)$ -retract, then it is permutably reducible or semilinear.*

LEMMA 3.3 (Case $2 < \kappa \leq n - 3$, [Kro08b]). *Let f be an n -quasigroup of arbitrary order and $\kappa(f) \in \{3, \dots, n - 3\}$. Then f is permutably reducible.* In [Kro08a], an example of irreducible n -quasigroup of order 4 whose $(n - 1)$ -retracts are all reducible is constructed for every even $n \geq 4$. So, the assumption of Lemma 3.3 cannot be extended to the case $\kappa = n - 2$. Nevertheless, in Section 5 we will prove the following:

LEMMA 3.4 (Case $\kappa = n - 2$). *Let $n \geq 5$. If an n -quasigroup f of order 4 has a semilinear permutably irreducible $(n - 2)$ -retract and all the $(n - 1)$ -retracts are permutably reducible, then f is permutably reducible or semilinear.*

The last case, announced in [Pot06], will be proved in Section 4:

LEMMA 3.5 (Case $\kappa = 2$). *Let $n \geq 5$; let f be an n -quasigroup of order 4, and let all its k -retracts with $2 < k < n$ be permutably reducible. Then f is permutably reducible.*

Proof of Theorem 3.1. The validity of the Theorem for $n \leq 4$ (and even for $n \leq 5$) is proved by exhaustion. Assume, by induction, that all m -quasigroups of order 4 with $m < n$ are reducible or semilinear. Consider an n -quasigroup f of order 4. It has

an irreducible $\kappa(f)$ -retract, which is semilinear by inductive assumption. Depending on the value of $\kappa(f) = 2, 3 \dots n-3, n-2, n-1$, the statement of Theorem follows from one of Lemmas 3.5, 3.3, 3.4, 3.2. $\square$

4. Proof of Lemma 3.5. We will prove a stronger variant (Lemma 4.2) of the statement. It uses the following concept:

DEFINITION 4.1. *An n -quasigroup f is called completely reducible if it is permutably reducible and all its principal retracts of arity more than 2 are permutably reducible (equivalently, f can be represented as a composition of $n-1$ binary quasigroups, e.g. $f(x_1, x_2, x_3, x_4, x_5) = f_1(f_2(x_1, x_3), f_3(f_4(x_2, x_5), x_4))$).*

LEMMA 4.2. *Let all the principal 3- and 4-retracts of an n -quasigroup f of order 4 ($n \geq 5$) be permutably reducible. Then f is completely reducible.*


DEFINITION 4.3. *An n -quasigroup f is called normalized if for all $i \in \{1, \dots, n\}$ and $a \in \Sigma$ it is true that $f(\bar{0}^{[i]}[a]) = a$.*

Denote by Γ the set of all (four) normalized binary quasigroups of order 4. It is straightforward that the operations from Γ are associative and commutative (they are isomorphic to the additive groups Z_2^2 and Z_4), and we will use the form $a \star b$ instead of $\star(a, b)$ to write the result of $\star \in \Gamma$.

Let $K_n = \langle V(K_n), E(K_n) \rangle$ be the complete graph with n vertices associated with the arguments $x_1, \dots, x_n$ of an n -ary operation. For the edges, we will use the short notation like $x_i x_j$. For any normalized n -quasigroup f we define the edge coloring $\mu_f : E(K_n) \rightarrow \Gamma$ by the following way: the color $\mu_f(x_i x_j)$ of an edge $x_i x_j \in E(K_n)$ is defined as the binary operation $\star$ such that $f(\bar{0}^{[i,j]}[x_i, x_j]) \equiv x_i \star x_j$.

PROPOSITION 4.4. *Let f be an n -quasigroup, $n \geq 5$, and let all 3- and 4-retracts of f be reducible. Then the coloring μ_f of K_n satisfies the following:*


- (A) *Every triangle is colored by at most 2 colors.*
- (B) *If a tetrahedron is colored by 2 colors with 3 edges of each color, then it includes a one-color triangle; i. e., the following fragment is forbidden:*



Proof. Every 3-retract of f is a composition of two binary operations, which yields (A).

Consider the 4-retract f_4 of f that corresponds to some four vertices (the other variables are fixed by 0). Since it is reducible and normalized, it can be represented as a composition of some normalized 2-quasigroup and 3-quasigroup. The 3-quasigroup is a 3-retract of f and, in its turn, can be represented as a composition of two normalized 2-quasigroups. So, $f_4(x, y, z, u)$, up to permutation of arguments, has the form $(x \star y) \circ (u \diamond v)$ or $x \circ (y \star (u \diamond v))$ for some $\star, \circ, \diamond \in \Gamma$. As follows from the hypothesis of (B), two of these three operations coincide. Thus, there are only four types of decomposition of f_4 : $(x \star y) \circ (u \star v)$, $x \star (y \circ u \circ v)$, $x \star y \star (u \circ v)$, $x \circ (y \star (u \circ v))$. In any case, (B) holds. $\square$

PROPOSITION 4.5. *Assume that an edge coloring μ of K_n satisfies (A) and (B). Then for each pairwise different $a, b, c, d \in V(K_n)$ the condition $\mu(ab) = \mu(ac) \neq \mu(bc) = \mu(bd) \neq \mu(cd)$ implies $\mu(ad) = \mu(ab)$. I. e.,*



Proof. Obviously, any other variant for $\mu(ad)$ contradicts (A) or (B). $\square$

The following proposition is easy to check.

PROPOSITION 4.6. *Let f and g be reducible 3-quasigroups of order 4, and let $f(x, y, 0) \equiv g(x, y, 0)$, $f(x, 0, z) \equiv g(x, 0, z)$, and $f(0, y, z) \equiv g(0, y, z)$. Then $f \equiv g$.*

Remark. Indeed, Proposition 4.6 holds for every order with the extra condition that f is a composition of two different or associative 2-quasigroups. The similar statement for n -quasigroups with $n > 3$ holds for an arbitrary order without extra conditions [KPS08, Theorem 1]: if two reducible n -quasigroups coincide on every n -tuples with one zero, then they are identical.

COROLLARY 4.7. *Let f and g be n -quasigroups of order 4 ($n \geq 3$) whose principal 3-retracts are all reducible. Assume that $f(\bar{0}^{[i,j]}[y, z]) \equiv g(\bar{0}^{[i,j]}[y, z])$ for every $i, j \in \{1, \dots, n\}$ and $y, z \in \Sigma$. Then $f \equiv g$.*

Proof. The equality $f(\bar{x}) \equiv g(\bar{x})$ is proved by induction on the number of non-zero elements in $\bar{x}$, using the reducibility of 3-retracts and Proposition 4.6.

For example, to prove that $f(1, 2, 3, 2, 1, \bar{0}) = g(1, 2, 3, 2, 1, \bar{0})$ we can consider the 3-retracts $f^3(x, y, z) \triangleq f(1, 2, x, y, z, \bar{0})$ and $g^3(x, y, z) \triangleq g(1, 2, x, y, z, \bar{0})$. By the induction assumption f^3 and g^3 meet the hypothesis of Proposition 4.6. Thus, $f^3 \equiv g^3$, and, in particular, $f^3(3, 2, 1) = g^3(3, 2, 1)$. $\square$

The following proposition is the key statement in the proof of Lemma 4.2.

PROPOSITION 4.8. *Assume that an edge coloring $\mu : E(K_n) \rightarrow \Gamma$ of the graph K_n meets (A) and (B). Then there exists a completely reducible n -quasigroup f such that $\mu_f = \mu$.*

Before proving Proposition 4.8 by induction, we consider one auxiliary statement, which will be used in the induction step. We say that an edge $xy \in E(K_n)$ is *inner* with respect to some edge coloring μ of K_n if for any $z \in V(K_n) \setminus \{x, y\}$ it is true that $\mu(xz) = \mu(yz)$.

PROPOSITION 4.9. *Assume that an edge coloring μ of K_n meets (A) and (B). Then K_n contains an inner edge.*

Proof. Consider an arbitrary sequence of edges $e_1, e_2, \dots, e_k$ that satisfies the following:

- (C) for every $j \in \{1, \dots, k-1\}$ the edges e_j and e_{j+1} are adjacent and $\mu(e_j) = \mu(e_j \triangle e_{j+1}) \neq \mu(e_{j+1})$ (where $\triangle$ means the symmetrical difference between two sets).

Denote by a_j the element from $e_j \setminus e_{j+1}$.




FIGURE 4.1. An example of a sequence $e_1, \dots, e_k$ from the proof of Proposition 4.8

Claim ().* We claim that for every i, j , $1 \leq i < j \leq k$, and $d \in e_j$ the vertices a_i and d are different and $\mu(a_i d) = \mu(e_i)$ (see Fig. 4.1). We will show this by induction on $j - i$.

If $j - i = 1$, the claim follows from (C). If $j - i = 2$, the claim follows from Proposition 4.5 ($a := a_i$, $b := a_{i+1}$). Assume $j - i > 2$. By the inductive assumption,

$$\begin{aligned} \mu(e_i) &= \mu(a_i a_{i+1}) = \mu(a_i a_{i+2}) \\ &\neq \mu(e_{i+1}) = \mu(a_{i+1} a_{i+2}) = \mu(a_{i+1} d) \\ &\neq \mu(e_{i+2}) = \mu(a_{i+2} d). \end{aligned}$$

Consequently, $a_i \neq d$, and, by Proposition 4.5 ($a := a_i$, $b := a_{i+1}$, $c := a_{i+2}$), we have $\mu(a_i d) = \mu(e_i)$. Claim (*) is proved.

So, all $a_1, \dots, a_{k-1}$ are mutually different, and thus there exists a maximum sequence $e_1, e_2, \dots, e_k$ satisfying (C). Then, its maximality and (*) imply that the edge e_k is inner. $\square$

Proof of Proposition 4.8. We will proceed by induction on n . If $n = 3$, then the statement is trivially true. Assume that Proposition 4.8 holds for $(n-1)$ -quasigroups. Consider an inner edge $e \in E(K_n)$. Without loss of generality we can assume $e = x_{n-1}x_n$, $\mu(e) = \star$. Denote by μ^{n-1} the restriction of the coloring μ on $K_{n-1} \subset K_n$. By the inductive assumption, there exists a completely reducible $(n-1)$ -quasigroup g such that $\mu_g = \mu^{n-1}$. Then $f(x_1, \dots, x_n) \triangleq g(x_1, x_2, \dots, x_{n-2}, x_{n-1} \star x_n)$ is a desired n -quasigroup (indeed, $\mu_f(x_{n-1}x_n) = \star = \mu(x_{n-1}x_n)$; if $i < j < n$, then $\mu_f(x_i x_j) = \mu_g(x_i x_j) = \mu(x_i x_j)$; if $i < n-1$, then $\mu_f(x_i x_n) = \mu_g(x_i x_{n-1}) = \mu(x_i x_{n-1}) = \mu(x_i x_n)$, where the last equality follows from the innerness of $x_{n-1}x_n$). $\square$

Proof of Lemma 4.2. Let f be an n -quasigroup of order 4 whose 3- and 4-retracts are all reducible. Without loss of generality we assume that f is normalized (otherwise, we can normalize it, applying an appropriate isotopy). Then, by Proposition 4.4, the corresponding edge coloring μ_f of the graph K_n satisfies (A) and (B). By Proposition 4.8, there exists a completely reducible n -quasigroup g with $\mu_g \equiv \mu_f$. By Corollary 4.7, f and g are identical. $\square$

5. Proof of Lemma 3.4. In the proof, we will use the following three propositions. The first simple one, on a representation of a reducible n -quasigroup with an irreducible $(n-1)$ -retract, holds for an arbitrary order.

PROPOSITION 5.1. *Assume that a reducible n -quasigroup D ($n \geq 3$) of an arbitrary order has an irreducible $(n-1)$ -retract $F\langle x_0, \dots, x_{n-1} \rangle \equiv D\langle x_0, \dots, x_{n-1}, 0 \rangle$. Then there are $i \in \{0, \dots, n\}$ and a 2-quasigroup h such that $h(x, 0) \equiv x$ and*

$$(5.1) \quad D\langle x_0, \dots, x_n \rangle \equiv F\langle x_0, \dots, x_{i-1}, h(x_i, x_n), x_{i+1}, \dots, x_{n-1} \rangle.$$

Proof. Since D is reducible, $D\langle x_0, \dots, x_n \rangle$ can be represented as $H\langle f(\bar{x}'), \bar{x}'' \rangle$ where $\bar{x}'$ and $\bar{x}''$ are disjoint groups of variables, each containing at least two variables. If x_n is grouped with more than one other variable, then fixing x_n gives a reducible retract, which contradicts the irreducibility of F . So, we conclude that for some $i \in \{0, \dots, n-1\}$ there exists one of the following two representations of D :

$$(5.2) \quad \begin{aligned} D\langle x_0, \dots, x_n \rangle &\equiv G\langle g(x_i, x_n), \tilde{x} \rangle, \\ D\langle x_0, \dots, x_n \rangle &\equiv g\langle G(\tilde{x}), x_i, x_n \rangle \end{aligned}$$

where $\tilde{x} \triangleq (x_0, \dots, x_{i-1}, x_{i+1}, \dots, x_{n-1})$ and G and g are $(n-1)$ - and 2- quasigroups. Moreover, the existence of a representation of the first type implies the existence of a representation of the second type, and vice versa:

$$g\langle G(\tilde{x}), x_i, x_n \rangle \equiv \begin{cases} 1 & \text{if } G(\tilde{x}) = g(x_i, x_n) \\ 0 & \text{if } G(\tilde{x}) \neq g(x_i, x_n) \end{cases} \equiv G\langle g(x_i, x_n), \tilde{x} \rangle.$$

So, we can assume that (5.2) holds. Put $\gamma(x_i) \triangleq g(x_i, 0)$. Then, $F\langle x_0, \dots, x_{n-1} \rangle \equiv G\langle \gamma(x_i), \bar{x} \rangle$, and (5.1) holds with $h(x_i, x_n) \triangleq \gamma^{-1}g(x_i, x_n)$. $\square$

In what follows, permutations $\sigma : \Sigma \rightarrow \Sigma$ will be denoted by the value lists $(\sigma(0), \sigma(1), \sigma(2), \sigma(3))$; denote $Id \triangleq (0, 1, 2, 3)$.

PROPOSITION 5.2 (on autotopies of a semilinear n -quasigroup). *Assume f is a standardly semilinear n -quasigroup. Denote by π the permutation $(1, 0, 3, 2)$. Then for every different $i, j \in \{0, \dots, n\}$ the following holds:*

- a) $f\langle \bar{x} \rangle \equiv f\langle \bar{x}^{[i,j]}[\pi x_i, \pi x_j] \rangle$, where $\bar{x} \triangleq (x_0, \dots, x_n)$;
- b) if $f\langle \bar{x} \rangle \equiv f\langle \bar{x}^{[i,j]}[\mu x_i, \nu x_j] \rangle$ holds for some other pair of non-identity permutations $(\mu, \nu) \neq (\pi, \pi)$ and $n \geq 3$, then f is reducible.

Proof. a) It is straightforward that $f\langle \bar{x}^{[i]}[\pi x_i] \rangle \equiv L\langle \bar{x} \rangle - f\langle \bar{x} \rangle$, where $L\langle \cdot \rangle$ is from Definition 2.5. So, $f\langle \bar{x}^{[i,j]}[\pi x_i, \pi x_j] \rangle \equiv L\langle \bar{x} \rangle - (L\langle \bar{x} \rangle - f\langle \bar{x} \rangle) \equiv f\langle \bar{x} \rangle$.

b) Without loss of generality assume that $i = 1, j = 2$. Put

$$\begin{aligned} \alpha(x, y) &\triangleq f(x, y, \bar{0}), \\ \beta(x, \bar{z}) &\triangleq f(x, 0, \bar{z}), \quad \bar{z} \triangleq (z_1, \dots, z_{n-2}), \\ \gamma(x) &\triangleq f(x, 0, \bar{0}). \end{aligned}$$

Assume there exists a pair (μ, ν) that satisfies the hypothesis of b). Then $\alpha(x, y) \equiv \alpha(\mu x, \nu y)$. It is easy to see that the permutation ν does not have fixed points. So, ν is either a cyclic permutation or an involution $((2, 3, 0, 1)$ or $(3, 2, 1, 0))$ different from $\pi = (1, 0, 3, 2)$. In any case, we can derive the following:

Claim ()*. For each $v \in \Sigma$ there exist permutations $\rho_v, \tau_v : \Sigma \rightarrow \Sigma$ such that $f(x, y, \bar{z}) \equiv f(\rho_v x, \tau_v y, \bar{z})$ and $\tau_v v = 0$ (in other words, the group of permutations τ admitting $f(x, y, \bar{z}) \equiv f(\rho x, \tau y, \bar{z})$ for some ρ acts transitively on Σ , i.e., has only one orbit):

Case 1. If ν is a cyclic permutation, then $v, \nu v, \nu^2 v, \nu^3 v$ are pairwise different; so, one of the pairs $(Id, Id), (\mu, \nu), (\mu^2, \nu^2), (\mu^3, \nu^3)$ can be chosen as (ρ_v, τ_v) , proving (*).

Case 2. If ν is $(2, 3, 0, 1)$ or $(3, 2, 1, 0)$, then $v, \nu v, \pi v, \nu \pi v$ are pairwise different, and (ρ_v, τ_v) can be chosen from $(Id, Id), (\mu, \nu), (\pi, \pi), (\mu \pi, \nu \pi)$.

Claim (*) is proved. Then,

$$\begin{aligned} f(x, y, \bar{z}) &\equiv f(\rho_y x, \tau_y y, \bar{z}) \equiv f(\rho_y x, 0, \bar{z}) \equiv \beta(\rho_y x, \bar{z}) \\ &\equiv \beta(\gamma^{-1} \alpha(\rho_y x, 0), \bar{z}) \equiv \beta(\gamma^{-1} \alpha(\rho_y x, \tau_y y), \bar{z}) \equiv \beta(\gamma^{-1} \alpha(x, y), \bar{z}) \end{aligned}$$

and thus f is reducible provided $n \geq 3$. $\square$

The next proposition concerns 2-quasigroups of order 4, and the proof is straightforward.

PROPOSITION 5.3. *Let s and t be 2-quasigroups. Denote $s_i(x) \triangleq s(x, i)$ and $t_i(x) \triangleq t(x, i)$. Let $s_0 = t_0 = Id$, and let for every i either $t_i s_i^{-1} = Id$ or $t_i s_i^{-1} = (1, 0, 3, 2)$. Then either $s \equiv t$ or for some permutation ϕ the 2-quasigroup $s'(x, y) \triangleq s(x, \phi y)$ is standardly semilinear.*

Proof. Denote $\pi \triangleq (1, 0, 3, 2)$; observe $\pi = \pi^{-1}$. Assume that $s \neq t$. Then there are at least two different elements $i, j \in \{1, 2, 3\}$ for which $t_i s_i^{-1} = \pi$. Denote by k the third element, i.e., $\{i, j, k\} = \{1, 2, 3\}$. The permutation t_i has no fixed points, otherwise there is a contradiction with t_0 ; similarly, $s_i = \pi t_i$ has no fixed points. So, t_i and, similarly, t_j belong to $\{(2, 3, 0, 1), (2, 3, 1, 0), (3, 2, 0, 1), (3, 2, 1, 0)\}$. The only variant for t_k is $(1, 0, 3, 2)$. Then, $s'(x, y) \triangleq s(x, \phi y)$ is standardly semilinear with $\phi \triangleq (0, k, i, j)$. $\square$

Proof of Lemma 3.4. Assume C is an n -quasigroup of order 4. Assume all the $(n-1)$ -retracts of C are reducible and C has a semilinear irreducible $(n-2)$ -retract E . Without loss of generality assume that

$$E\langle x_0, \dots, x_{n-2} \rangle \equiv C\langle x_0, \dots, x_{n-2}, 0, 0 \rangle$$

and E is standardly semilinear. We will use the following notation for retracts of C (the table illustrates their mutual arrangement, where the last and last-but-one coordinates of Σ^{n+1} are thought as ordinate and abscissa respectively; for example, B_2 corresponds to fixing abscissa by 2):

$$\begin{aligned} E_{a,b}\langle x_0, \dots, x_{n-2} \rangle &\triangleq C\langle x_0, \dots, x_{n-2}, a, b \rangle, \\ A_b\langle x_0, \dots, x_{n-2}, y \rangle &\triangleq C\langle x_0, \dots, x_{n-2}, y, b \rangle, \\ B_a\langle x_0, \dots, x_{n-2}, z \rangle &\triangleq C\langle x_0, \dots, x_{n-2}, a, z \rangle. \end{aligned}$$

Since A_0 is reducible and fixing $y := 0$ leads to the irreducible E , by Proposition 5.1 we have

$$(5.3) \quad A_0\langle x_0, \dots, x_{n-2}, y \rangle \equiv E\langle x_0, \dots, x_{i-1}, h(x_i, y), x_{i+1}, \dots, x_{n-2} \rangle$$

for some $i \in \{0, \dots, n-2\}$ and 2-quasigroup h such that $h(x_i, 0) \equiv x_i$.

From (5.3), we see that all the retracts $E_{a,0}$, $a \in \Sigma$, are isotopic to E . Similarly, we can get the following:

Claim ().* All the retracts $E_{a,b}$, $a, b \in \Sigma$ are isotopic to E .

Then, we conclude that a representation similar to (5.3) is valid for every $b \in \Sigma$:

$$A_b\langle x_0, \dots, x_{n-2}, y \rangle \equiv E_{0,b}\langle x_0, \dots, x_{i_b-1}, h_b(x_{i_b}, y), x_{i_b+1}, \dots, x_{n-2} \rangle$$

for some $i \in \{0, \dots, n-2\}$ and 2-quasigroup h_b such that $h_b(x, 0) \equiv x$.

*Claim (**).* We claim that i_b does not depend on b . Indeed, assume, for example, that $i_1 = 0$ and $i_2 = 1$, i. e.,

$$\begin{aligned} A_1\langle x_0, \dots, x_{n-2}, y \rangle &\equiv E_{0,1}\langle h_1(x_0, y), x_1, x_2, \dots, x_{n-2} \rangle, \\ A_2\langle x_0, \dots, x_{n-2}, y \rangle &\equiv E_{0,2}\langle x_0, h_2(x_1, y), x_2, \dots, x_{n-2} \rangle. \end{aligned}$$

Then, fixing x_0 in the first case leads to a retract isotopic to E ; fixing x_0 in the second case leads to a reducible retract (recall that $n \geq 5$). But, analogously to (*), these two retracts are isotopic; this contradicts the irreducibility of E and proves (**).

Without loss of generality we can assume that $i_b = 0$, i. e.,

$$(5.4) \quad A_b\langle x_0, x_1, \tilde{x}_2, y \rangle \equiv E_{0,b}\langle h_b(x_0, y), x_1, \tilde{x}_2 \rangle;$$

here and later $\tilde{x}_2 \triangleq (x_2, \dots, x_{n-2})$. Similarly, we can assume without loss of generality that either

$$(5.5) \quad B_a\langle x_0, x_1, \tilde{x}_2, z \rangle \equiv E_{a,0}\langle g_a(x_0, z), x_1, \tilde{x}_2 \rangle$$

or

$$(5.6) \quad B_a\langle x_0, x_1, \tilde{x}_2, z \rangle \equiv E_{a,0}\langle x_0, g_a(x_1, z), \tilde{x}_2 \rangle$$

where 2-quasigroups g_a satisfy $g_a(x, 0) \equiv x$.

Using (5.4) and (5.5), we derive

$$\begin{aligned}
 (5.7) \quad C\langle x_0, x_1, \tilde{x}_2, y, z \rangle &\equiv A_z\langle x_0, x_1, \tilde{x}_2, y \rangle \\
 &\equiv E_{0,z}\langle h_z(x_0, y), x_1, \tilde{x}_2 \rangle \\
 &\equiv B_0\langle h_z(x_0, y), x_1, \tilde{x}_2, z \rangle \\
 &\equiv E_{0,0}\langle g_0(h_z(x_0, y), z), x_1, \tilde{x}_2 \rangle,
 \end{aligned}$$

which means that C is reducible, because $f(x, y, z) \triangleq g_0(h_z(x, y), z)$ must be a 3-quasigroup. So, it remains to consider the case (5.6). Consider two subcases.

Case 1. The 2-quasigroup g_a does not depend on a ; denote $g \triangleq g_a$. Then, repeating the first three steps of (5.7) and applying (5.6), we derive that

$$C\langle x_0, x_1, \tilde{x}_2, y, z \rangle \equiv E_{0,0}\langle h_0(x_0, y), g(x_1, z), \tilde{x}_2 \rangle,$$

and C is reducible.

Case 2. For some fixed a we have $g_0 \neq g_a$; denote $s_i(x) \triangleq g_0(x, i)$, $t_i(x) \triangleq g_a(x, i)$, and $r_i(x) \triangleq h_i(x, a)$. From (5.4), we see that

$$(5.8) \quad E_{a,0}\langle x_0, x_1, \tilde{x}_2 \rangle \equiv E_{0,0}\langle r_0(x_0), x_1, \tilde{x}_2 \rangle,$$

$$(5.9) \quad E_{a,b}\langle x_0, x_1, \tilde{x}_2 \rangle \equiv E_{0,b}\langle r_b(x_0), x_1, \tilde{x}_2 \rangle.$$

From (5.6), we see that

$$(5.10) \quad E_{0,b}\langle x_0, x_1, \tilde{x}_2 \rangle \equiv E_{0,0}\langle x_0, s_b(x_1), \tilde{x}_2 \rangle,$$

$$(5.11) \quad E_{a,b}\langle x_0, x_1, \tilde{x}_2 \rangle \equiv E_{a,0}\langle x_0, t_b(x_1), \tilde{x}_2 \rangle.$$

Applying consecutively (5.10), (5.9), (5.11), and (5.8), we find that for each b the retract $E = E_{0,0}$ satisfies

$$E\langle x_0, x_1, \tilde{x}_2 \rangle \equiv E_{0,b}\langle \dots \rangle \equiv E_{a,b}\langle \dots \rangle \equiv E_{a,0}\langle \dots \rangle \equiv E\langle r_0 r_b^{-1} x_0, t_b s_b^{-1} x_1, \tilde{x}_2 \rangle.$$

By Proposition 5.2, the irreducibility of E means that $t_b s_b^{-1} \in \{Id, (1,0,3,2)\}$ for every b . By Proposition 5.3, for some permutation ϕ the 2-quasigroup $s(x, z) \triangleq g_0(x, \phi z)$ is standardly semilinear. Since a composition of standardly semilinear quasigroups is a standardly semilinear quasigroup, we see that B_0 is a semilinear $(n-1)$ -quasigroup. Lemma 3.2 completes the proof. $\square$

6. Acknowledgement. The authors thank the referees for their work in reviewing the manuscript and the audience of the seminar “Coding Theory” in the Sobolev Institute of Mathematics for the patience during reporting this result.

REFERENCES

- [AHS04] S. V. AVGUSTINOVICH, O. HEDEN, AND F. I. SOLOV'YEV, *The classification of some perfect codes*, Des. Codes Cryptography, 31(3) (2004), 313–318.
DOI: 10.1023/B:DESI.0000015891.01562.c1
- [Bel72] V. D. BELOUSOV, *n-Ary Quasigroups*, Shtiintsa, Kishinev, 1972. In Russian.
- [Kro08a] D. S. KROTOV, *On irreducible n-ary quasigroups with reducible retracts*, Eur. J. Comb., 29(2) (2008), 507–513.
DOI: 10.1016/j.ejc.2007.01.005
ArXiv: math/0607785

- [Kro08b] D. S. KROTOV, *On reducibility of n -ary quasigroups*, Discrete Math., 308(22) (2008), 5289–5297.
DOI: 10.1016/j.disc.2007.08.099
ArXiv: math/0607284
- [KPS08] D. S. KROTOV, V. N. POTAPOV, AND P. V. SOKOLOVA, *On reconstructing reducible n -ary quasigroups and switching subquasigroups*, Quasigroups Relat. Syst., 16(1) (2008), 55–67.
ArXiv: math/0608269
- [LM98] C. F. LAYWINE AND G. L. MULLEN, *Discrete Mathematics Using Latin Squares*, Wiley, New York, 1998.
- [Pot06] V. N. POTAPOV, *On completely commutatively reducible n -quasigroups*, in Proceedings of the XVI International School-Seminar “Synthesis and Complexity of Controlling Systems”, St.-Petersburg, Russia, June 2006, 88–91. In Russian.
- [PK06] V. N. POTAPOV AND D. S. KROTOV, *Asymptotics for the number of n -quasigroups of order 4*, Sib. Math. J., 47(4) (2006), 720–731. Translated from Sib. Mat. Zh. 47(4) (2006), 873–887.
DOI: 10.1007/s11202-006-0083-9
ArXiv: math/0605104

FURTHER READING

- [KP] D. S. KROTOV AND V. N. POTAPOV, *On reducibility of n -ary quasigroups, II*.
ArXiv: 0801.0055